

Anticoagulation Day to Day Usage - Quick Reference Guide

This quick reference guide explains how to use the **Anticoagulation Manager** template for monitoring patients on Warfarin. It is a training aid and as such is not exhaustive. For more information, please refer to the online help <u>https://help.cegedim-</u>

healthcare.co.uk/Visionplus/Content/G_Full_Help_Topics/Outcomes_Manag er/Anti-Coagulation/main.htm.

Accessing the Anticoagulation Manager Clinical Template

This section explains how to view and add data to the **Anticoagulation Manager** template.

- 1. From Consultation Manager, select a patient and then either:
 - From the Windows Notification Area, right click Vision+ and select Clinical Templates Anticoagulation Manager:



Highlight the Anticoagulation Manager category and select
 Template from the toolbar:



2. The **Anticoagulation Pathway** tab displays, unless the patient has a possible alcohol-related interaction, in which case it shows the **Warfarin** tab with the interaction highlighted.





3. To add data, complete any relevant information by selecting **Codeset** at the end of the appropriate row and ticking **Apply** against the clinical code required:

ome Appearance		
	Mon 06/08/2018 -	
e & Codeset Show Ind	icator 🌱 Previous Tab 🥐 Next Tab	
le	Controls	
ry Anticoagulation Pathway Warfarin	R DOACs Other Assessment	
	Anticoagulation	Codeset
vision	Managor	
A Cegedim Brand	Mallagel	1 I
On anticoagulation pathway	No Data Recorded	V UIII
Anticoagulant Therapy		
Anticoagulant Therapy		
Anticoagulant Therapy To indicate treatment has stopped use	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets are the patient is monitored by a District nurse	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse	wise use 66Q5. 🛛 👻 🛄 🎔 📷
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP Comments	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP Comments	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP Comments	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped use (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP Comments Please record either the planned dural life-long anticoagulation therapy.	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse tion of treatment or the planned stop date in the 'Comments' box, or if the	wise use 66Q5.
Anticoagulant Therapy To indicate treatment has stopped usi (Warfarin therapy stopped) to indicate Prescribed anticoagulants Please record instances below whe Monitoring by District nurse Shared care: district nurse/GP Comments Please record either the planned durat life-long anticoagulation therapy. Planned duration of treatment	e the read code 66c1. (Anticoagulant therapy stopped) for DOACs other warfarin treatment has stopped 10/01/2018: Warfarin 500microgram tablets ere the patient is monitored by a District nurse tion of treatment or the planned stop date in the 'Comments' box, or if the 05/05/2018: Duration of anticoagulant therapy	wise use 66Q5.

Note - When adding data, you can select a different date, priority or add freetext comments, if appropriate.

- 4. After all relevant information has been recorded on the **Anticoagulation Manager Pathway** tab, select the **Warfarin** tab.
- 5. From here you can review and record the following:

and the second	
Adheres to medication regimen O Current medictn reported by p	
Comments	

- If the patient adheres to their medication, tick **Adheres to their medication regimen**.
- That the patient has reported their current medication.



Using the Integrated Warfarin Dosing Tool

This section explains how to use the dosing calculator:

- 1. From the Warfarin tab, highlight Warfarin Dose (using the integrated dosing tool) and select Warfarin Dose .
- 2. The **Warfarin dose** screen displays the patient's current dosing history at the top. The patient's INR results populates from the electronic lab results, if not, please record it manually in the **New INR and Dose** section below:

Warfarin dose								
🚡 Settings 🛛 💆 P	rint Pre <u>v</u> iew	<u>P</u> rint	Print A4 Re	port				
INR & Dose Treatm	ent & Advice His	story Tren	ids					
Current INR and	Dose							
Start Date 25/06/2018	Indication Calf vein	o n thrombus				INR tar 2.00-3.00	get D [2.50]	
INR Date 04/07/2018	INR 3.00	←	INR result					
Dose Date	Wed	Thu	Fri	Sat	Sun	Mon	Tue	Total Dose (mg)
04/07/2018	3.00	3.00	3.00	3.00	3.00	3.00	3.00	21.0
Recall Date 11/07/2018								

3. The Advised Warfarin Adjustment displays on the bottom half of the screen. If correct, make sure there is no tick in Same Daily Dose and select Use:

New INR and Do	ose								
INR Date 06/08/2018	INR 4	Cle	ar					Select L	Use
Advised Warfarin 19.1 mg/week	Adjustment [Coventry / W	√arwickshir	e]	O Rour	nd Up (19.5)	Round	Down (19.0)	
Same Daily Dose	■ ←	No tick				200 - 000			
Dose Date 06/08/2018	Mon	Tue	Wed	Thu	Fri	Sat	Sun	Total Dose (mg)	6
Recall Date									
	•								
									_
								OK Canc	:el

Note - The **Same Daily Dose** option is there for elderly patients who may struggle with their dose varying from day to day, and should be used in exceptional cases and not as the standard practice.

4. For doses not achieved with the tablet strengths available, the option to round up or round down the weekly dose displays with the recommended option selected.



- 5. The weekly dosing regime displays, and can be printed and given to the patient. Select either **Print A4 Report** or to print the details on the right-hand side of a prescription, select **Print**.
- 6. The recommended recall is also calculated based on the patient's INR history.

Training Tip - The recommended recall date does not record in the patient's record. If you want to enter this information, select the INR tab on the Anticoagulation Manager template and select Next planned INR test date.

- 7. Select **OK** to save the changes and exit the **Warfarin dose** screen.
- 8. To record that the patient was advised about the new Warfarin dose, select the INR tab and select Codeset III on the Patient informed of latest INR result entry:

INR result	06/08/2018: International normalised ratio	4	× 🛄 🕋
Target INR	06/08/2018: Warfarin upper target	3.0	¥ 🛄 💌
TTR calculator	06/08/2018: International normalised ratio	4	* 🔚 🛄 🖎
Patient informed of latest INR result	No Data Recorded		× 🛄 📭
Next planned INR test date			

- 9. The **Codeset Codes** screen displays, tick **Apply** against the code required and select **OK**.
- 10.Select **Save and Close** to record the data into the patient record.